

Objective Daily Maps of Thermocline Depth for the SYNOP Eastern Array

Z. R. Hallock Oceanography Division Ocean Science Directorate



Approved for public release; distribution is unlimited. Naval Oceanographic and Atmospheric Research Laboratory, Stennis Space Center, Mississippi 39529-5004.

ABSTRACT

A major component of the field activities for the northwest Atlantic Regional Energetics Experiment (REX) is focused on the acquisition and analysis of data from inverted echo sounders (IESs). This technical note describes an analysis of data records from nine IESs deployed in an array near 40° N, 55°W. IES acoustic travel-time data were converted to thermocline depths, which were in turn objectively interpolated and contoured, resulting in daily maps. These maps provide a useful tool for studying the evolution of mesoscale features in the area of the experiment.



Acces	on For	
DTIC	ounced	N
By Distrib	ution /	
•	wailability C	Codes
Dist	Avail and Special	/ or
A-1		

Acknowledgments

Thanks are extended to Dr. Amy Bower of the Woods Hole Oceanographic Institution for providing the correlation function used in the analysis. This work was supported by the Office of Naval Research under Program Element 0601153N, Dr. H. C. Eppert, Jr., Program Manager, as part of the basic research project "Extended GEOSAT Mission."

Contents

Introduction	1
Conversion of IES travel times to Z12	1
Objective mapping of Z12	2
References	54

OBJECTIVE DAILY MAPS OF THERMOCLINE DEPTH FOR THE SYNOP EASTERN ARRAY

Introduction

The final phase of NOARL's (Naval Oceanographic and Atmospheric Research Laboratory) Regional Energetics Experiment (REX) coincided with the beginning of ONR's (Office of Naval Research) SYNOP (Synoptic Ocean Prediction) program, which was conducted between Cape Hatteras and the Grand Banks in the Gulf Stream region. The SYNOP field experiments extended from late 1987 through most of 1990 and consisted of extensive moored and hydrographic measurements of oceanographic parameters in three regions: the Inlet Array (near Cape Hatteras), the Central Array (near 68°W) and the Eastern Array (near 55°W and further east). The REX component consisted chiefly of an array of 13 inverted echo sounders (IESs) deployed in a pattern to complement SYNOP current meter moorings in the eastern array. Figure 1 shows the location of the IESs in perspective, and Figure 2 shows the details of IES and current meter mooring positions. Superimposed on Figures 1 and 2 are GEOSAT-ERM groundtracks. The IES data (as well as CTD and XBT data) acquired during this experiment are described by Teague et al. (1990). IES logistics are listed in Table 1; four of the instruments were not recovered.

The purpose of the REX field programs (as well as one of the main objectives of SYNOP) was to monitor fluctuations of the Gulf Stream and to describe its variability and dynamics. Results from REX field efforts are described by Hallock et al. (1989, 1991) and Teague and Hallock (1990).

In this technical note we describe the conversion of time series of IES-measured acoustic travel times to time series of thermocline depth (in particular, Z12, the depth of the 12° isotherm) and, subsequently, objective mapping of the Z12 data over the SYNOP Eastern Array, producing a series of daily maps of Z12. The daily maps thus produced appear following the description of the method.

Conversion of IES travel times to Z12

The REX-SYNOP IESs in the Eastern Array yielded 9 records of approximately 260 days of acoustic travel time (tau). These data were low-pass filtered (with half-amplitude point at 40 hrs) to remove tidal and higher frequencies. Changes in tau (tau') about record means were converted to changes in Z12 (Z12') using regression coefficients based on historical CTD data from the region of the Eastern Array. To obtain Z12 from Z12' a reference point for each record was established using "calibration" XBTs launched at the IES position just prior to recovery (Watts and Johns, 1982). In particular, with CTD data from the Eastern Array region in summer 1983 (Hogg et al., 1985) and in early spring 1988 (Hall and Fofonoff, 1991) regression of calculated Z12' on calculated tau' yielded:

 $Z12=23.1(tau'-tau_0')+Z12_0$

where tau', tau_0 ' are in milliseconds, tau_0 ' is tau' at the end of the record and Z12 $_0$ is Z12 from calibration XBTs. The overall uncertainty in the resulting Z12 is about 30 m (this is essentially the error-of-fit of the CTD-based tau'-Z12' regressions). Table 2 lists basic statistics of the Z12 time series.

Objective mapping of Z12

Objective analysis (OA) (Bretherton et al., 1976; Watts et al., 1989) was used to interpolate the IES Z12 data to produce a series of daily maps of Z12 for the Eastern Array. In this method, the original observations, Z_j (j=1,9 here), at points X_j , are interpolated to points X_k to form a regular grid that can be contoured. This interpolation is effected with a weighting function, A_{ik} , such that

$$Y_k = \sum_j A_{jk} Y_j \tag{1}$$

In the OA method the weighting function is based on the covariances of the variable to be interpolated and is given by

$$A_{ik} = \sum_{j} C_{ij}^{-1} C_{jk}$$
 (2)

where $C_{ij} = C(Y_i, Y_j)$ is the normalized covariance matrix. An estimate of the normalized error in variance associated with each interpolated point can be calculated

$$e_k = 1 - \sum_i \sum_j A_{ik} A_{jk} C_{ij}. \qquad (3)$$

In this analysis, the covariance function, C, is assumed to be homogeneous and isotropic (dependent only on the distance between points), i.e., $C(X_i, X_j) = C(lX_i - X_j l) = C(r_{ij})$. This assumption may not always be justified in the vicinity of features such as the Gulf Stream, as pointed out by Hogg (SYNOPtician 2(6), December 1991).

The OA first required the removal of a mean field and normalization by a standard deviation (sd) field. Since the NOARL-IES spatial distribution was quite sparse and the instruments were deployed for only about 9 months, the mean and sd fields suggested by Bower and Hogg (SYNOPtician 2(3), June 1991; henceforth BH) were used; these were based on over 2 years of observations in the Eastern Array.

Since the BH functions are for temperature at 500 m (not Z12) they were scaled in amplitude prior to use in the analysis. This was done by least squares fitting the BH functions,

mean = a - b tanh
$$(\frac{y - 50}{100.8})$$

sd = c exp $[-(\frac{y - 50}{259})^2]$ (4)

to the sample Z12 means and sd's (those appearing in table 2). The values of the coefficients were found to be:

$$a = 453 \text{ m}$$

 $b = 336 \text{ m}$
 $c = 133 \text{ m}$.

For the same reasons given above, a correlation function developed by BH was also used. Correlations calculated with the actual Eastern Array IES records showed considerable scatter but bracketed the function used by BH:

$$C(r) = S_0 \exp\left[-\left(\frac{r}{S_2}\right)^2\right] \cos(2\pi r/S_1)$$
 (5)

where

$$S_0 = 0.902$$

 $S_1 = 610 \text{ km}$
 $S_2 = 141 \text{ km}$

(Bower, personal communication).

A 10×13 grid (dx=dy=25 km) was selected and daily maps of Z12 were generated by calculating Y_k (eq. 1) for each grid point. The Y_k were then redimensionalized by multiplying by the sd and adding back the mean (eq. 4). Actual mean and sd fields were calculated with the gridded data and contoured and appear as figures 3 and 4. Figure 5 is the set of daily maps of Z12 starting on page 11. The error field is superimposed on each of the daily maps. The error contours are in increments of 10%; the dotted contour near the center of each map is 10%, the dashed contour is 20% and the remaining solid contours are 30%, 40%, . . ., etc. Beyond 30% the results become questionable.

Table 1. Times and locations of the IES/PGs.

IES No.	Latitude (N)	Longitude (W)	Depth (m)	Deployed	Recovered		
IES87001	34°23.2	55°03.2	5454	13 Sept 87	27 June 88		
IES87002	38°58.5	53°55.4	5412	14 Sept 87	28 June 88		
IES87003	38°58.9	54°40.3	5393	13 Sept 87	27 June 88		
IES87004	38°59.0	55°24.1	5409	13 Sept 87	26 June 88		
IES87005	39°33.8	55°46.0	5369	15 Sept 87	26 June 88		
IES87006	39°33.7	55°01.1	5367	14 Sept 87	30 June 88		
IES87007	39°33.2	54°18.1	5351	14 Sept 87	29 June 88		
IES87008	Not Recovered						
IES87009	Not Recovered						
IES87010	Not Recovered						
IES87011	40°07.3	55°23.5	5324	15 Sept 87	5 July 88		
IES87012	40°41.0	55°01.7	5213	16 Sept 87	4 July 88		
IES87013	Not Recovered			•			

Table 2. Statistics of Z12 series.

IES No.	Min.	Max.	Mean	Sd.
87-001	459.	859.	671.	81.
87-002	455.	882.	718.	67.
87-003	362.	921.	762.	67.
87-004	447.	866.	730.	72.
87-005	200.	860.	642.	105.
87-006	272.	901.	630.	143.
87-007	177.	832.	563.	192.
87-011	63.	806.	403.	184.
87-012	-147.	522.	191.	131.




Figure 1. Location of NRL IES array deployed from September 1987 to June 1988. GEOSAT groundtracks are superimposed.




Figure 2. Detail of IES locations (circles; open circles indicate instruments not recovered) and Woods Hole current meter mooring locations (triangles). GEOSAT ground tracks are superimposed.




Figure 3. Map of mean Z12 calculated using the objectively interpolated, gridded data.








Figure 4. Map of standard deviation of Z12 about the mean shown in Figure 3.


Figure 5. Objective maps of Z12 in SYNOP Eastern Array for 29 September 1987 to 11 June 1988. Percent error variance contours superimposed on Z12 maps: dotted-10%, dashed-20%, heavy solid-30% and higher. IES positions marked by +.





























































































































References

- Bretherton, P., R. E. Davis, and C. B. Fandry (1976). A Technique for Objective Analysis and Design of Oceanographic Experiments Applied to MODE-73. *Deep Sea Res.*, 23, 559-582.
- Hall, M. M. and N. P. Fofonoff (1991). Downstream Development of the Gulf Stream from 68° W to 55° W. J. Phys. Oceanogr., submitted.
- Hallock, Z. R., J. L. Mitchell, and J.D. Thompson (1989). Sea Surface Height Variability near the New England Seamounts: an Intercomparison among in situ Observations, GEOSAT Altimetry and Numerical Simulations. J. Geophys. Res. C4 94(66), 8021-8028.
- Hallock, Z. R., W. J. Teague, and J. D. Thompson (1991). A Comparison of Observed and Modeled Sea-surface Topographic Time-series near the New England Seamounts. J. Geophys. Res., 96, C7, 12,635-12,644.
- Hogg, N. G., R. S. Pickart, R. M. Hendry, and W. Smethie (1985). On the Northern Recirculation Gyre of the Gulf Stream. *Deep Sea Res.*, 33(9A), 1139-1166.
- Teague, W. J., and Z. R. Hallock (1990). Gulf Stream Path Analysis near the New England Seamounts. J. Geophys. Res., 95 C2, 1647-1662.
- Watts, D. R., K. L. Tracey, and A. I. Friedlander (1989). Producing Accurate Maps of the Gulf Stream Thermal Front using Objective Analysis. J. Geophys. Res. 94, 8040-8052.
- Watts, D. R. and W. E. Johns (1982). Gulf Stream Meanders: Observations on Propagation and Growth. J. Geophys. Res., 87(C2), 9467-9476.

Distribution List

David Adamec Laboratory for Oceans Goddard Space Flight Center Greenbelt, MD 20770

John Bane Univ. of North Carolina Curriculum of Marine Sciences Venable Hall 045-A Chapel Hill, NC 27514

Amy Bower Physical Oceanography Clark Building 315A WHOI Woods Hole, MA 02543

Dale Bretschneider ONR Physical Oceanography Program Code 1122PO Arlington, VA 22217-5000

Everett Carter Naval Postgraduate School OC/CR Monterey, CA 93940

Dr. Alan Chave AT&T IE444 600 Mountain Avenue Murray Hill, NJ 07974

Peter Cornillon Narragansett Bay Campus Rm. 112 - Watkins Bldg. Narragansett, RI 02882

Meghan Cronin Narragansett Bay Campus Rm. 113 - Watkins Bldg. Narragansett, RI 02882

David Evans
ONR
Physical Oceanography Program
Code 1122ML
Arlington, VA 22217-5000

Erik Fields Narragansett Bay Campus Rm. 219 - Watkins Bldg. Narragansett, RI 02882 Charles N. Flagg Brookhaven National Lab Bldg. 218 Upton, NY 11713

Glenn R. Flierl
Department of Meteorology
M.I.T.
Rm. 54-1422
Cambridge, MA 02139

Dr. N. P. Fofonoff Physical Oceanography Dept. Clark Building 3 WHOI Woods Hole, MA 02543

Sandra Fontana Narragansett Bay Campus Rm. 221 - Watkins Bldg. Narragansett, RI 02882

Avijit Gangopahdyay Dept. Earth & Planetary Science Pierce Hall Harvard University Cambridge, MA 02138

Craig Gilman Narragansett Bay Campus Rm. 211 - Watkins Bldg. Narragansett, RI 02882

Dr. Dale Haidvogel
Institute of Marine & Coastal
Sciences
Rutgers University
P. O. Box 231
New Brunswick, NJ 08903-0231

Melinda Hall Physical Oceanography Clark Building 3 WHOI Woods Hole, MA 02543

George Halliwell RSMAS University of Miami 4600 Rickenbacker Causeway Miami, FL 33149 Ross Hendry Atlantic Oceanography Lab Bedford Institute of Oceanography Dartmouth, Nova Scotia CANADA B2Y 4A2

Nelson Hogg WHOI Woods Hole, MA 02543

William Holland NCAR P. O. Box 3000 Boulder, CO 80307

Steve Howden Narragansett Bay Campus Rm. 219 - Watkins Bldg. Narragansett, RI 02882

Rui Xin Huang Dept. of Physical Oceanography Clark Building 3 WHOI Woods Hole, MA 02543

William Johns RSMAS 4600 Rickenbacker Causeway Miami, FL 33149

Kathie Kelly Clark Building 3 WHOI Woods Hole, MA 02543

HyunSook Kim Narragansett Bay Campus Rm. 113 - Watkins Bldg. Narragansett, RI 02882

Thomas Kinder
Office of Naval Research
Physical Oceanography Program
Code 1121CS
Arlington, VA 22217-5000

Dr. Peter Koltermann WOCE - International Program Ofc. 105 Deacon Lab - Brook Rd. Wormley, Godalming Surrey, GU85UB UNITED KINGDOM Harris Kontoyiannis Narragansett Bay Campus Rm. 219 - Watkins Bldg. Narragansett, RI 02882

Richard Lambert Physical Oceanography Program National Science Foundation 1800 G Street, NW Washington, DC 20550

Chris Langdon Lamont-Doherty Geological Observatory Columbia University Palisades, NY 10964

John Lillibridge 4726 Boiling Brook Parkway Rockville, MD 20852

Steven Lohrenz USM/CMS Bldg. T-242 SSC, MS 39529

Susan Lozier Physical Oceanography Department Clark Building 2 WHOI Woods Hole, MA 20543

Dr. Douglas Luther Scripps Institute of Oceanography UCSD Code A-030 La Jolla, CA 92093

Prof. Paola Malanotte-Rizzoli Dept. of Earth & Atmos. Plan. Science MIT 54-1416 Cambridge, MA 02139

Arthur Mariano RSMAS/MPO University of Miami 4600 Rickenbacker Causeway Miami, FL 33149

George Mellor Princeton University Pheasant Hill Road Princeton, NJ 08540 Prof. Chris Mooers RSMAS University of Miami 4600 Rickenbacker Causeway Miami, FL 33149

Donald B. Olson RSMAS/MPO University of Miami 4600 Rickenbacker Causeway Miami, FL 33149

Peter B. Ortner NOAA, OAR/ERL/AOML/OCD Room 609 4501 Rickenbacker Causeway Miami, FL 33149

J. L. Pelegri Oceanography Department Old Dominion University Norfolk, VA 23529-0276

Louise Perkins
Department of Earth Atm. Plan.
Science
MIT 54-1420
Cambridge, MA 02139

Robert Pickart Clark Building 3 WHOI Woods Hole, MA 20543

Richard Pieper University of Southern California University Park Los Angeles, CA 90089

Lawrence Pratt
Physical Oceanography Department
Clark Building 3
WHOI
Woods Hole, MA 02543

Allan Robinson Harvard University 29 Oxford St. Cambridge, MA 02138

Tom Rossby Narragansett Bay Campus Rm. 222 - Watkins Bldg. Narragansett, RI 02882 William Schmitz, Jr. WHOI Clark Building 3 Woods Hole, MA 02543

Thomas Shay University of North Carolina Curriculum of Marine Sciences Venable Hall 045-A Chapel Hill, NC 27514

Tao Song Narragansett Bay Campus Rm. 221 - Watkins Bldg. Narragansett, RI 02882

Michael Spall WHOI Clark Building 3 Woods Hole, MA 02543

Thomas Spence Division of Ocean Sciences National Science Foundation 1800 G Street, NW Washington, DC 20550

Kathy Schultz-Tokos Institut für Meerskunde Düsternbrooker Weg 20 D2300 Kiel 1 . FEDERAL REPUBLIC OF GERMANY

Dr. Jorge Vasquez Jet Propulsion Lab Mail Stop 169-236 4800 Oak Drive Pasadena, CA 91205

Randy Watts Narragansett Bay Campus Rm. 212 - Watkins Building Narragansett, RI 02882

Charles Yentsch Bigelow Laboratory McKown Point W. Boothbay Harbor, ME 04575

Norm Zabusky
Dept. of Mech. and Aerospace Eng.
Rutgers University/Busch Campus
P. O. Box 909
Piscataway, NJ 08855-0909

```
Naval Research Laboratory
Stennis Space Center, MS 39529-5004
Att: Code 331
      Code 330
      Code 320
      Code 300
      Code 210 (H. Morris)
      Code 323 (D. Thompson)
      Code 323 (J. Mitchell)
      Code 323 (H. Hurlburt)
      Code 331 (W. Teague)
      Code 331 (D. Johnson)
      Code 125L (10)
      Code 125P
Naval Research Laboratory
Washington, DC 20375
Att: Commanding Officer
      Library (2)
Naval Oceanographic Office
Stennis Space Center, MS 39522-5001
Att: Commanding Officer
      Code TD
      Library (2)
      John Blaha (TDT)
      Charles Horton (OSTM)
```

Vance Sprague (OP)

Form Approved REPORT DOCUMENTATION PAGE OBM No. 0704-0188 Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202 4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503. 1. Agency Use Only (Leave blank). 2. Report Date. 3. Report Type and Dates Covered. May 1992 Final 4. Title and Subtitle. 5. Funding Numbers. Program Element No. 0601153N Objective Daily Maps of Thermocline Depth for the SYNOP Eastern Array Project No. 03208 Author(s). Task No. 3CG Z.R. Hallock Accession No. DN394458 Work Unit No. 13311V Performing Organization Name(s) and Address(es). 8. Performing Organization Report Number. Naval Oceanographic and Atmospheric Research Laboratory Ocean Science Directorate NOARL Technical Note 259 Stennis Space Center, MS 39529-5004 9. Sponsoring/Monitoring Agency Name(s) and Address(es). 10. Sponsoring/Monitoring Agency Report Number. Naval Oceanographic and Atmospheric Research Laboratory **Basic Research Management Office NOARL Technical Note 259** Stennis Space Center, MS 39529-5004 11. Supplementary Notes. 12a. Distribution/Availability Statement. 12b. Distribution Code. Approved for public release; distribution is unlimited. 13. Abstract (Maximum 200 words). A major component of the field activities for the northwest Atlantic Regional Energetics Experiment (REX) is focused on the acquisition and analysis of data from inverted echo sounders (IESs). This technical note describes an analysis of data records from nine IESs deployed in an array near 40°N, 55°W. IES acoustic travel-time data were converted to thermocline depths, which were in turn objectively interpolated and contoured, resulting in daily maps. These maps provide a useful tool for studying the evolution of mesoscale features in the area of the experiment. 14. Subject Terms. 15. Number of Pages.

of Report.

17. Security Classification

Altimetry, Mesoscale Oceanography, Ocean Forecasting

18. Security Classification

of This Page.

Unclassified

20. Limitation of Abstract.

57

19. Security Classification

of Abstract.

Unclassified

16. Price Code.

SAR